NORTH FORK VETERINARY CLINIC

12080 Locksley Ln. Auburn, CA 95602 530-888-8788

Puppy/K-9 Vaccine Protocols:

Vaccine Vaccine Interval

DHPP-Parvo (5 in 1) Multiple doses given at 6, 9, 12 and 16 weeks

• If started at 8 wks, then 8, 12, 16 weeks

Rabies Single 1 year duration given at 16 weeks

Lepto (4 serovar) Two doses given at 12 and 16 weeks

For at risk dogs...exposed to wildlife/forest/acreage/hiking trails

Not recommended in small dogs/sensitive breeds

Two doses given at 12 and 16 weeks Lyme

• From tick bites

Rattlesnake Two doses given at 16 and 20 weeks (or older)

For at risk dogs...exposed to rural areas, hiking and/or forest environments

Second vaccine given 4 weeks apart

Bordetella (Kennel Cough) Single (intranasal) dose given at 6 weeks or older (***required to board at NFVC)

Single dose with a yearly booster

Given to at risk dogs...exposed to boarding kennels/dog parks/dog shows/obedience

school/agility classes

Influenza (H3N8/H3N2) Two doses given after 7 weeks of age and then booster 2 weeks later (***required to board at NFVC)

Given to at risk dogs...exposed to boarding kennels/dog parks/dog shows/obedience

school/agility classes

Adult K-9 Vaccine Protocols:

Vaccine Vaccine Interval

DHPP-Parvo (5 in 1) Booster given at 1 year, then every 3 years

• If history is unknown, then 2 doses given 4 weeks apart

Rabies Booster given at 1 year, then every 3 years

• If history is unknown, then single dose w/booster at 1 yr.

Lepto (4 serovar) Single dose given yearly

Best before rainy season

• If it's the first time the vaccine has been given, then 2 doses given 4 weeks apart is recommended

Lyme Single dose given yearly

• If it's the first time the vaccine has been given, then 2 doses given 4 weeks apart is recommended

Rattlesnake Single dose given yearly

Best given in the spring.

If it's the first time the vaccine has been given, then 2 doses given 4 weeks apart is recommended

Single (intranasal) dose given yearly (***required to board at NFVC) Bordetella (Kennel Cough)

Influenza (H3N8/H3N2) Two doses give at least 2 weeks apart, then booster annually (***required to board at NFVC)

> Given to at risk dogs...exposed to boarding kennels/dog parks/dog shows/obedience school/agility classes

Vaccine Risk Assessment:

Do	g's Name:		Last Name:	Date:		
1.	Current Age:	Breed		Male/Female (circle)		
2.	How did you acquire own	nership? (circle)				
	 From a pet store From a breeder From an adoption Stray or from a residual 	n agency				
3.	List any other localities other than the greater Auburn/Foothill area that your pet has lived:					
4.	Do you travel with your p	et? Where?				
5.	5. Does your dog live in a household with multiple pets? Y / N (circle)					
6.	6. Is your dog exposed to: horses / livestock / fowl (circle all that apply)					
7.	7. Is your dog around children or seniors on a typical day? Y / N (circle)					
8.	8. Does anyone in your dog's environment have an impaired immune system? Y / N (circle)					
9. Is your dog outdoors more that 50% of the time? Y / N (circle)						
10	 Stay in a fenced Go on leash wal Visit dog parks Visit public area Stay in a neighb Go camping or least of the company of the compan	yard ks s orhood only niking trails (where these with stagnant pone facilities accilities classes/dog shows	here is wildlife)			
11	Date of last / previous vaccinations, heartworm test:					
12	Has your dog ever ha	d a reaction to vacci	ines? Y / N (circle) Wh	ich one?		

Other considerations: heartworm area /flea-tick risk/GI parasites/worms

13.